

# Decreased Anterior Cingulate Volume in Combat-Related PTSD

Steven H. Woodward, Danny G. Kaloupek, Chris C. Streeter, Christelle Martinez, Marie Schaer, and Stephan Eliez

**Background:** Neuroanatomical data point to functional relationships between the anterior cingulate cortex (ACC) and subcortical centers regulating fear, in particular, the amygdala. Functional brain imaging has disclosed divergent patterns of ACC activation in persons with posttraumatic stress disorder (PTSD). In addition, two preliminary structural imaging studies have found evidence of smaller ACC volume in PTSD. We explored associations between PTSD and ACC volume in a relatively large sample of adult combat veterans in which PTSD, lifetime alcohol abuse/dependence, and Vietnam versus Gulf War service were crossed.

**Methods:** Subjects were US military combat veterans of the Vietnam and Gulf Wars recruited from two metropolitan areas served by allied Department of Veterans Affairs PTSD treatment/research centers. Anterior cingulate cortex volume was analyzed as a function of grouping factors with and without adjustment for body size.

**Results:** Posttraumatic stress disorder was associated with smaller anterior cingulate cortex volume. This effect persisted in subjects without histories of alcoholism, did not interact with cohort effects, and was not modified by adjustment for body size.

**Conclusions:** Anterior cingulate cortex volume is substantially smaller in association with combat-related PTSD, a finding broadly consistent with cingulate hypofunctionality in that disorder.

**Key Words:** Stress disorders, posttraumatic, magnetic resonance imaging, gyrus cinguli

Keen interest has emerged in the role that the anterior cingulate cortex (ACC) might play in posttraumatic stress disorder (PTSD) (Hamner et al 1999; Pitman et al 2001; Taber et al 2003; Villarreal and King 2001). Of particular interest are this region's close neuroanatomical relationships with subcortical components of the "central fear system," including the amygdala and locus coeruleus (Carmichael and Price 1995; Jodo and Aston-Jones 1997; Jodo et al 1998; McDonald et al 1996; Vogt et al 1979, 1987, 1995; Vogt and Pandya 1987). If the ACC supplies inhibitory regulation of the amygdala, then attenuation of that regulation could provide an avenue for understanding multiple features of PTSD. Absolute or relative hypoactivation of ACC during tasks involving exposure to traumatic reminders in persons with PTSD has now been replicated in three laboratories (Bremner et al 1999a, 1999b; Lanius et al 2001, 2003; Shin et al 1999, 2001). Accompanying such findings are indications that the ACC is coupled to the hypothalamic-pituitary-adrenal (HPA) axis. The cingulate cortex contains the largest concentration of receptors for corticotropin-releasing factor (CRF) among all cortical areas in the brain of the rhesus macaque (Sanchez et al 1999). As well, there have been observations of pyramidal cell apical dendritic remodeling in ACC following glucocorticoid challenge (Wellman 2001) and restraint stress (Radley et al 2004), similar to those originally motivating exploration of hippocampal volume in PTSD (Bremner et al 1995; Pitman et al 2001). Taken together,

these findings suggest that reduced ACC volume might be found in association with PTSD.

Two preliminary structural magnetic resonance imaging (MRI) studies have made observations compatible with this hypothesis. Rauch et al (2003) reported smaller pregenual ACC and subcallosal cortex volumes (the latter combining portions of Brodmann's areas 32, 11, and 25) in nine Vietnam combat nurses with PTSD as compared with nine combat nurse control subjects. These authors did not find smaller dorsal ACC volume and invoked the preferential coupling of pregenual ACC and subgenual cortex to amygdala to explain this specificity. In their sample, ACC volume reductions were not associated with comorbid major depression. Using voxel-based morphometry, Yamasue et al (2003) found evidence of ACC gray matter hypodensity in 9 survivors of the Tokyo subway sarin attack with diagnoses of lifetime PTSD (1 also met criteria for current PTSD) as compared with 16 survivors who had never met criteria for PTSD. Here, we report on an effort to replicate these findings in a well-characterized sample of 99 survivors of combat and military operational stress. This sample also allowed us to account for variance associated with lifetime alcoholism and, to a limited degree, with normal aging.

## Methods and Materials

### Recruitment and Screening

Subjects provided written informed consent in accordance with procedures of the Institutional Review Boards of either Stanford University Medical School/VA Palo Alto Healthcare System or Boston VA Medical Center and the McLean Hospital. Subjects were recruited through a combination of advertising and word-of-mouth contacts with current and past patients and research volunteers. Initial screening established that subjects were US military veterans of the Vietnam Conflict or the Persian Gulf War (GW) who had been exposed to substantial military operational stress but now reported no current or past central nervous system (CNS) disease or psychosis and no alcohol or substance abuse/dependence in the last 6 months. Screening exclusions were based on low stress exposure, current alcohol or substance use, high fever(s), loss of consciousness requiring medical attention, or known contraindications to magnetic reso-

From the Clinical Laboratory and Education Division and Psychology Service (SHW), National Center for PTSD, VA Palo Alto Healthcare System, Palo Alto, California; Behavioral Science Division (DGK), National Center for PTSD, VA Boston Healthcare System, Division of Psychiatry (DGK, CCS), Boston University School of Medicine, Boston, Massachusetts; and Department of Psychiatry (CM, MS, SE), University of Geneva, Geneva, Switzerland.

Address reprint requests to Steven H. Woodward, Ph.D., National Center for PTSD, Clinical Laboratory and Education Division, Mail Code 334 PTSD, VA Palo Alto HCS, 3801 Miranda Avenue, Palo Alto, CA 94034; E-mail: steve.woodward@med.va.gov

Received January 3, 2005; revised July 8, 2005; accepted July 27, 2005.

nance (MR) scanning. As a result of formal diagnosis, additional volunteers were excluded because they were negative for current military PTSD but positive for lifetime civilian PTSD (18), or were positive for current/recent alcohol/drug abuse (14), probable brain damage (6), or psychosis (2). In addition, four subjects later withdrew due to fatigue or nicotine withdrawal; two missed their scanning appointments and were unreachable; and five withdrew due to claustrophobia. After participating, 11 additional subjects were excluded due to imaging artifact and 2 due to previously undiagnosed brain injury. The final sample included 55 subjects from the Palo Alto site and 44 from the Boston site. Completing subjects were paid \$200.

Final sample characteristics are presented in Table 1. The sample included 99 military veterans of either the Vietnam Conflict or the Gulf War. Subjects classified as PTSD-positive (PTSD+) met criteria for current PTSD as a result of experiencing one or more military traumas. Subjects classified as PTSD-negative (PTSD-) had also been exposed to military operational stress but were free of diagnosable PTSD, current or lifetime, due to military or civilian trauma. (All control subjects met DSM-IV PTSD Criterion A1, and all but three met Criterion A2). Alcohol abuse/dependence-positive (ETOH+) subjects were so classified based on meeting lifetime but not current alcohol abuse or dependence criteria (minimum period since meeting criteria: 6 months). Gulf War veterans had a mean age of 38 years, and Vietnam veterans had a mean age of 56 years.

### Psychometrics

Subjects meeting screening criteria were administered both interview and self-report measures. Semistructured interviewing was conducted by trained and experienced Masters-level staff using the Clinician-Administered PTSD Scale (CAPS) (Blake et al 1997) for PTSD symptoms and selected Axis I modules of the Structured Clinical Interview for the DSM-IV (SCID) (First et al 1995) addressing mood episodes (e.g., major depression), psy-

chotic and associated symptoms, alcohol and other substance use disorders, and anxiety and other disorders (e.g., panic disorder). Self-report instruments included the Combat Exposure Scale (CES) (Keane et al 1989), the Life Events Checklist (LEC) (Blake et al 2000), the Mississippi Scale for Combat-Related PTSD (MISS) (Keane et al 1988), the Beck Depression Inventory (BDI) (Beck et al 1961), and the Michigan Alcohol Screening Test-Short Form (SMAST) (Selzer 1971).

### Brain Imaging

Magnetic resonance imaging was performed using two 1.5 T General Electric Signa scanners (GE, Fairfield, Connecticut) at similar hardware and software revisions, one at the Diagnostic Radiology Center of Veterans Affairs Palo Alto Healthcare System and one at the Brain Imaging Center of McLean Hospital (Belmont, Massachusetts). Coronal images were acquired with a three-dimensional (3-D) volumetric pulse sequence (repetition time [TR] = 35 milliseconds, echo time [TE] = 6 milliseconds, flip angle = 45°, number of excitations [NEX] = 1, matrix size = 256 × 192, field of view = 24 cm<sup>2</sup>, slice thickness = 1.5–1.7 mm, 124 slices). Image optimization was performed in BrainImage (BrainImage 5.x, Stanford University, Stanford, California) following the standard protocols of the Stanford Psychiatry Neuroimaging Laboratory. Image optimizations included correction for inhomogeneity artifact, resampling to cubic (.9375 mm<sup>3</sup>) voxels, positional normalization by reference to the anterior and posterior commissures and intrahemispheric fissure, skull-stripping, tissue segmentation based on a constrained fuzzy algorithm (Reiss et al 1998), and parcellation according to a modified Talairach grid (Kates et al 1999; Talairach and Tournoux 1988).


Manual delineation of the cingulate cortex followed a protocol developed by one of the authors (SE). Left and right cingulate gyri were first traced in sagittal view on slices 5 millimeters lateral to the midline. Next, a coronal view was used to draw the superior, inferior, lateral, and medial boundaries of the cingulate

**Table 1.** Subject Characteristics and Anterior Cingulate Cortex Volumes

	Vietnam Cohort		Persian Gulf Cohort		PTSD	COHORT	ETOH	Interactions
	PTSD+	PTSD–	PTSD+	PTSD–				
N	38	25	13	23				
Age	53.5 (2.6)	56.0 (3.5)	37.0 (5.7)	36.7 (3.9)	n.s.	$p < .001$	$p = .011$	PTSD × COHORT $p = .027$ PTSD × COHORT × ETOH $p = .001$
% Male	100	100	77	83	n.s.	$p < .001$	n.s.	n.s.
% Caucasian	65.8	92.0	53.8	69.6	$p = .041$	n.s.	$p = .041$	n.s.
% Current MDD	78.9	4.0	69.2	4.3	$p < .001$	$p = .037$	n.s.	n.s.
% Lifetime MDD	89.5	28.0	76.9	17.4	$p < .001$	$p = .012$	n.s.	n.s.
% Lifetime ETOH	44.7	44.0	46.2	39.1	n.s.	n.s.		n.s.
Age at ETOH + Onset	24.6 (8.0)	21.5 (8.7)	26.7 (5.0)	22.1 (5.9)	n.s.	n.s.		n.s.
Years of Education	14.4 (1.8)	15.5 (2.2)	14.3 (1.7)	15.0 (1.9)	$p = .016$	n.s.	n.s.	n.s.
Combat Exposure Scale	29.8 (9.9)	24.2 (8.2)	19.9 (11.8)	8.6 (6.0)	$p < .001$	$p < .001$	n.s.	COHORT × ETOH $p = .041$
BDI	25.0 (8.9)	4.6 (3.7)	21.0 (7.3)	4.3 (4.0)	$p < .001$	n.s.	n.s.	n.s.
SMAST	3.9 (4.0)	2.1 (3.8)	3.3 (3.6)	0.5 (.9)	$p < .001$	$p = .035$	$p < .001$	PTSD × ETOH $p = .012$
MISS	122.8 (18.8)	68.2 (15.8)	107.8 (15.8)	59.0 (11.1)	$p < .001$	$p = .001$	$p = .038$	n.s.
CAPS Total Severity	75.9 (18.4)	8.8 (9.0)	75.9 (19.9)	8.4 (11.0)	$p < .001$	n.s.	n.s.	n.s.
WAIS Vocabulary Score	47.4 (12.0)	55.5 (7.1)	45.6 (12.4)	52.5 (8.0)	$p < .001$	n.s.	n.s.	PTSD × ETOH $p = .032$
Left ACC Volume in mL	6.9 (1.7)	8.6 (1.7)	6.6 (2.0)	7.6 (1.9)	$p < .001$	n.s.	n.s.	n.s.
Right ACC Volume	7.9 (3.0)	8.9 (2.0)	8.3 (2.1)	9.4 (2.1)				

Tabulation of demographic, diagnostic, and psychometric data by PTSD diagnosis and COHORT.  $p$ -values of main effects associated with the grouping factors are indicated in columns headed PTSD, COHORT, and ETOH, respectively. Column headed Interactions lists any interactions.

PTSD, posttraumatic stress disorder; ETOH, alcohol abuse/dependence; MDD, major depressive disorder; BDI, Beck Depression Inventory; SMAST, Michigan Alcohol Screening Test-Short Form; MISS, Mississippi Scale for Combat-Related PTSD; CAPS, Clinician-Administered PTSD Scale; WAIS, Wechsler Adult Intelligence Scale; ACC, anterior cingulate cortex; n.s., not significant.



cortex and adjacent white matter (see Figure 1). These boundaries were defined medially by the interhemispheric cortical surface and laterally by a line connecting the deepest extension of the cingulate sulcus to the deepest extent of cingulate gray matter ribbon directly superior to the corpus callosum (CC). The region of interest excluded tissue subjacent to the genu of the CC and posterior to a line dropped from the apex of the genu. Excessive intersubject variability in landmarks precluded delineation of subgenual cingulate cortex.) Posterior to the division of the splenium of the CC, the inferior border of the cingulate was defined by the calcarine fissure. Lastly, a dynamic Talairach grid was fitted on each brain (see Figure 2). In the rostrocaudal direction on a coronal anterior commissure-posterior commissure (AC-PC) oriented stack, Talairach sectors corresponding to B, C, D, and E1 in Figure 2 defined the anterior portion of the cingulate, while sectors E2, E3, F, G, and H defined the posterior portion. Two of the authors (C.M. and M.S.) performed all manual tracing of the cingulate blind to subject identity and diagnosis. Intraclass correlation coefficients of .94 calculated over two independent raters indicated good interrater reliability for cingulate volume measurements.

### Statistics

Primary analyses involved repeated-measures analysis of variance (ANOVA) with three grouping factors (PTSD, ETOH, COHORT) and two within factors (anterior versus posterior and right versus left). Follow-up analyses considered Talairach ACC subvolumes B through E1. Analyses of unadjusted volumes were repeated with adjustment for cerebral tissue volume, cranial volume, or Wechsler Adult Intelligence Scale (WAIS) vocabulary. Selected analyses were restricted to nonalcoholic subjects. Post-traumatic stress disorder effect sizes were estimated via “pooled  $d+$ ” (Hedges and Olkin 1985) to accommodate unequal numbers of GW and Vietnam cohort members across the four PTSD by ETOH subgroupings. Finally, correlational analyses were conducted on ACC volumes and selected continuous PTSD severity measures to assess parametric covariations.


### Results

The ACC was larger in the right hemisphere while the posterior cingulate cortex (PCC) was larger on the left [interaction of hemisphere and anterior-posterior dimension:  $F(1,91) = 25.6$ ,  $p < .001$ ]. There was no main effect of hemisphere and no interaction between hemisphere and any grouping factor. The ACC [ $F(1,91) = 11.7$ ,  $p = .001$ ] but not the PCC [ $F(1,91) = .056$ , n.s.] was smaller in subjects diagnosed with PTSD [interaction of PTSD and anterior-posterior factor:  $F(1,91) = 5.28$ ,  $p = .024$ ; see Figure 3]. The effect of PTSD on ACC volume did not interact with ETOH or with COHORT (all  $F$ s  $< 1$ ). Neither ETOH nor COHORT was associated with a main effect on ACC volume (all  $F$ s  $< 1$ ). No effect or interaction involving PTSD, ETOH, or COHORT on ACC volume was substantially modified by adjusting for cranial volume, cerebral tissue volume, or WAIS vocabulary. Likewise, the exclusion of seven female subjects had no impact on the results.

When repeated in ETOH- subjects, the effect of PTSD on total ACC volume persisted [ $F(1,51) = 6.77$ ,  $p = .012$ ]. Posttraumatic stress disorder effect sizes across the ETOH+ and ETOH- subsamples were comparable (ETOH+:  $d+ = -.71$ , 95% confi-

**Figure 1.** Successive coronal slices indicating manual tracings of cingulate cortical tissue areas on a typical brain.






**Figure 2.** Sagittal view of the cingulate region of interest (ROI) of a typical brain upon which has been superimposed a Talairach grid in accordance with standard locators.


dence interval [CI] =  $-1.34$  to  $-.09$ ; ETOH:  $d+ = -.78$ , 95% CI =  $-1.35$  to  $-.21$ ).

A second repeated-measures ANOVA was performed in which the within factors represented a finer parcellation of the ACC (two hemispheres by four Talairach ACC subvolumes B through E1). This analysis did not find a significant interaction of PTSD and subvolume [ $F(3,89) = 2.67$ ,  $p_{H-F} = .053$ ;  $\epsilon = .714$ ], though there was a trend toward a slightly larger effect at the B subvolume roughly corresponding to pregenual ACC (see Figure 4).

When computed across all subjects, a pattern of inverse correlations was evident between continuous measures of PTSD severity and ACC volume (CAPS total severity score:  $\rho[97] = -.332$ ,  $p < .001$ ; MISS:  $\rho[96] = -.339$ ,  $p < .001$ ); however, the correlation between ACC volume and BDI only approached significance ( $\rho[95] = -.197$ ,  $p < .053$ ).



**Figure 3.** Comparison of PTSD effects at anterior and posterior cingulate cortices. PTSD, posttraumatic stress disorder; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex.



**Figure 4.** Comparison of PTSD effects at Talairach subvolumes B–E1 of anterior cingulate cortex. PTSD, posttraumatic stress disorder.

## Discussion

This analysis of 99 combat veterans confirmed and extended previous findings. Anterior cingulate cortex volume was observed to be smaller in persons with combat-related PTSD whether or not they were diagnosed with lifetime alcoholism or belonged to one of two military/aging cohorts. Statistical adjustment for body size did not modify the results. We did not find that the effect of PTSD was limited to pregenual ACC. Our data suggested instead that dorsal ACC, often associated with more cognitive functions (Bush et al 2000), is also reduced in volume in PTSD. We did not assess subcallosal cortical volume and therefore we could not replicate this aspect of the observations of Rauch et al (2003).

To our knowledge, ACC volume has not been specifically considered in prior studies of the effects of alcoholism on the brain. The absence of a COHORT effect on ACC volume is compatible with two studies that failed to find normative aging effects on this structure (Jernigan et al 1991; Raz et al 1997). Acknowledging that the cohorts did not differ greatly in age, this absence is interesting in light of the exaggerated vulnerability to aging exhibited by nearby dorsolateral frontal cortex (Raz et al 1998, 2004).

The findings of this study are compatible with data suggesting the ACC is hypofunctional in PTSD. In addition to those studies noted in the introduction, functional imaging studies in normal humans using both MR and positron emission tomography (PET) have shown that the ACC is activated by tasks requiring behavioral monitoring/inhibition (Kerns et al 2004; Menon et al 2001) and emotion regulation (Ochsner et al 2002), both domains relevant to the clinical symptomatology of PTSD. Of special interest, Hariri et al (2003) found evidence of ACC involvement in the downregulation of amygdala responses to generic threat cues in normal subjects. The directionalities of all these results are compatible with the observed terminations of cortical afferents on inhibitory interneurons within the amygdala (Carmichael and Price 1995; McDonald et al 1996). Notwithstanding the appeal of a simple ACC hypoactivation model, *hyperactivation* of ACC in PTSD subjects during the generation of trauma-related images has also been reported (Lanius et al 2004; Rauch et al 1996; Shin et al 1997). Both Zubieta et al (1999) and Liberzon et al (1999) observed ACC hyperactivation in PTSD patients during exposure

to trauma-related sounds. Anterior cingulate cortex hyperactivation has also been observed during script-driven dissociation (Lanius et al 2002). Relatedly, Gilboa et al (2004) performed an analysis of interregional cerebral blood flow (CBF) correlations during script-driven imagery and found correlations between amygdala and ACC metabolism to be *positive*. As more studies become available, some of these inconsistencies may be resolved by consideration of differences among anterior cingulate subregions (Bush et al 2000; Vogt et al 1992), between tasks, or between the measurements employed. Until then, we should consider the full complement of functional interactions between anterior cingulate and amygdala to be potentially relevant to PTSD.

The effort to understand the contribution of ACC to PTSD can capitalize on a rich cognitive electrophysiological literature focusing on this region. Anterior cingulate cortex gives rise to multiple scalp-recordable electrophysiological events, including the error-related and feedback negativities (Herrmann et al 2004; Holroyd et al 1998; Mathalon et al 2003; Miltner et al 2003) and midline frontal theta (Asada et al 1999; Gevins et al 1997; Ishii et al 1999). These phenomena have already exhibited associations with anxiety (Aftanas and Golocheikine 2001; Hajcak et al 2003; Inanaga 1998). Their examination in PTSD is now strongly warranted.

*Major funding for this work was provided under a Veterans Affairs/Department of Defense (VA/DoD) assistance agreement to SHW and DGK from the US Army Medical Research and Materiel Command (DoD) administered through the Institute for Medical Research (VA). Special thanks are due to Stephen Grate and Lori Moll who facilitated the necessary arrangements at those institutions. Support was also provided by a National Institute on Alcohol Abuse and Alcoholism award (K23AA13149) and a VA Career Development Award to CCS. Support for SE was provided by the Swiss National Fund for Research Career Development Award (32-63135-00 and PP00B-102864/1).*

*We thank the Research Services of the VA Palo Alto and VA Boston Healthcare Systems for their assistance. Important institutional and technical support was also provided by the following individuals: Wendy K. Stegman, Lorraine P. Stewart, John E. Drace, Patricia Spezia, Carla Ambriz, and Gary E. Gold at the VA Palo Alto Healthcare System; Kelly Teresi, Frederick Kanter, Steven Blank, Susan Proctor, and Erica Stone at the VA Boston Healthcare System; Thomas J. Brosnan, Anil Patwardhan, and Eric Schmitt at Stanford University; Rosemond Villafuerte, Ann Smith, and Eileen Connolly at McLean Hospital; and Michael A. Dove at the Defense Manpower Database Command. We are also grateful for the early guidance provided by Kelvin O. Lim.*

- Aftanas LI, Golocheikine SA (2001): Human anterior and frontal midline theta and lower alpha reflect emotionally positive state and internalized attention: High-resolution EEG investigation of meditation. *Neurosci Lett* 310:57–60.
- Asada H, Fukuda Y, Tsunoda S, Yamaguchi M, Tonoike M (1999): Frontal midline theta rhythms reflect alternative activation of prefrontal cortex and anterior cingulate cortex in humans. *Neurosci Lett* 274:29–32.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J (1961): An inventory for measuring depression. *Arch Gen Psychiatry* 4:561–571.
- Blake D, Weathers F, Nagy L, Kaloupek D, Klauminzer G, Charney D, et al (2000): *Clinician-Administered PTSD Scale (CAPS) Instructional Manual*. Springfield, VA: National Center for PTSD, National Technical Information Service.
- Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Charney DS, Keane TM (1997): *Clinician-Administered PTSD Scale for DSM-IV: Current and Life-*

- time Version*. Springfield, VA: National Center for PTSD, National Technical Information Service.
- Bremner JD, Narayan M, Staib LH, Southwick SM, McGlashan T, Charney DS (1999a): Neural correlates of memories of childhood sexual abuse in women with and without posttraumatic stress disorder. *Am J Psychiatry* 156:1787–1795.
- Bremner JD, Randall P, Scott TM, Bronen RA, Seibyl JP, Southwick SM, et al (1995): MRI-based measurement of hippocampal volume in patients with combat-related posttraumatic stress disorder. *Am J Psychiatry* 152: 973–981.
- Bremner JD, Staib LH, Kaloupek D, Southwick SM, Soufer R, Charney DS (1999b): Neural correlates of exposure to traumatic pictures and sound in Vietnam combat veterans with and without posttraumatic stress disorder: A positron emission tomography study. *Biol Psychiatry* 45:806–816.
- Bush G, Luu P, Posner MI (2000): Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci* 4:215–222.
- Carmichael ST, Price JL (1995): Limbic connections of the orbital and medial prefrontal cortex in macaque monkeys. *J Comp Neurol* 363:615–641.
- First MB, Spitzer RL, Gibbon M, Williams JBW (1995): *Structured Clinical Interview for the DSM-IV Axis-I Disorders*. New York: Biometrics Research Department, New York State Psychiatric Institute.
- Gevins A, Smith ME, McEvoy L, Yu D (1997): High-resolution EEG mapping of cortical activation related to working memory: Effects of task difficulty, type of processing, and practice. *Cereb Cortex* 7:374–385.
- Gilboa A, Shalev AY, Laor L, Lester H, Louzoun Y, Chisin R, et al (2004): Functional connectivity of the prefrontal cortex and the amygdala in posttraumatic stress disorder. *Biol Psychiatry* 55:263–272.
- Hajcak G, McDonald N, Simons RF (2003): Anxiety and error-related brain activity. *Biol Psychol* 64:77–90.
- Hamner MB, Lorberbaum JP, George MS (1999): Potential role of the anterior cingulate cortex in PTSD: Review and hypothesis. *Depress Anxiety* 9:1–14.
- Hariri AR, Mattay VS, Tessitore A, Fera F, Weinberger DR (2003): Neocortical modulation of the amygdala response to fearful stimuli. *Biol Psychiatry* 53:494–501.
- Hedges LV, Olkin I (1985): *Statistical Methods for Meta-Analysis*. New York: Academic Press, Inc.
- Herrmann MJ, Rommler J, Ehls AC, Heidrich A, Fallgatter AJ (2004): Source localization (LORETA) of the error-related-negativity (ERN/Ne) and positivity (Pe). *Brain Res Cogn Brain Res* 20:294–299.
- Holroyd CB, Dien J, Coles MG (1998): Error-related scalp potentials elicited by hand and foot movements: Evidence for an output-independent error-processing system in humans. *Neurosci Lett* 242:65–68.
- Inanaga K (1998): Frontal midline theta rhythm and mental activity. *Psychiatry Clin Neurosci* 52:555–566.
- Ishii R, Shinosaki K, Ukai S, Inouye T, Ishihara T, Yoshimine T, et al (1999): Medial prefrontal cortex generates frontal midline theta rhythm. *Neuroreport* 10:675–679.
- Jernigan TL, Archibald SL, Berhow MT, Sowell ER, Foster DS, Hesselink JR (1991): Cerebral structure on MRI, Part I: Localization of age-related changes. *Biol Psychiatry* 29:55–67.
- Jodo E, Aston-Jones G (1997): Activation of locus coeruleus by prefrontal cortex is mediated by excitatory amino acid inputs. *Brain Res* 768:327–332.
- Jodo E, Chiang C, Aston-Jones G (1998): Potent excitatory influence of prefrontal cortex activity on noradrenergic locus coeruleus neurons. *Neuroscience* 83:63–79.
- Kates WR, Warsofsky IS, Patwardhan A, Abrams MT, Liu AM, Naidu S, et al (1999): Automated Talairach atlas-based parcellation and measurement of cerebral lobes in children. *Psychiatry Res* 91:11–30.
- Keane TM, Caddell JM, Taylor KL (1988): Mississippi Scale for Combat-Related Posttraumatic Stress Disorder: Three studies in reliability and validity. *J Consult Clin Psychol* 56:85–90.
- Keane TM, Fairbank JA, Caddell JM, Zimering RT, Taylor KL, Mora CA (1989): Clinical evaluation of a measure to assess combat exposure. *Psychol Assess* 1:53–55.
- Kerns JG, Cohen JD, MacDonald AW 3rd, Cho RY, Stenger VA, Carter CS (2004): Anterior cingulate conflict monitoring and adjustments in control. *Science* 303:1023–1026.
- Lanius RA, Williamson PC, Boksman K, Densmore M, Gupta M, Neufeld RW, et al (2002): Brain activation during script-driven imagery induced dissociation.

- tive responses in PTSD: A functional magnetic resonance imaging investigation. *Biol Psychiatry* 52:305–311.
- Lanius RA, Williamson PC, Densmore M, Boksman K, Gupta MA, Neufeld RW, et al (2001): Neural correlates of traumatic memories in posttraumatic stress disorder: A functional MRI investigation. *Am J Psychiatry* 158: 1920–1922.
- Lanius RA, Williamson PC, Densmore M, Boksman K, Neufeld RW, Gati JS, et al (2004): The nature of traumatic memories: A 4-T fMRI functional connectivity analysis. *Am J Psychiatry* 161:36–44.
- Lanius RA, Williamson PC, Hopper J, Densmore M, Boksman K, Gupta MA, et al (2003): Recall of emotional states in posttraumatic stress disorder: An fMRI investigation. *Biol Psychiatry* 53:204–210.
- Liberzon I, Taylor SF, Amdur R, Jung TD, Chamberlain KR, Minoshima S, et al (1999): Brain activation in PTSD in response to trauma-related stimuli. *Biol Psychiatry* 45:817–826.
- Mathalon DH, Whitfield SL, Ford JM (2003): Anatomy of an error: ERP and fMRI. *Biol Psychol* 64:119–141.
- McDonald AJ, Mascagni F, Guo L (1996): Projections of the medial and lateral prefrontal cortices to the amygdala: A Phaseolus vulgaris leucoagglutinin study in the rat. *Neuroscience* 71:55–75.
- Menon V, Adelman NE, White CD, Glover GH, Reiss AL (2001): Error-related brain activation during a Go/NoGo response inhibition task. *Hum Brain Mapp* 12:131–143.
- Miltner WH, Lemke U, Weiss T, Holroyd C, Scheffers MK, Coles MG (2003): Implementation of error-processing in the human anterior cingulate cortex: A source analysis of the magnetic equivalent of the error-related negativity. *Biol Psychol* 64:157–166.
- Ochsner KN, Bunge SA, Gross JJ, Gabrieli JD (2002): Rethinking feelings: An fMRI study of the cognitive regulation of emotion. *J Cogn Neurosci* 14: 1215–1229.
- Pitman RK, Shin LM, Rauch SL (2001): Investigating the pathogenesis of post-traumatic stress disorder with neuroimaging. *J Clin Psychiatry* 62:47–54.
- Radley JJ, Sisti HM, Hao J, Rocher AB, McCall T, Hof PR, et al (2004): Chronic behavioral stress induces apical dendritic reorganization in pyramidal neurons of the medial prefrontal cortex. *Neuroscience* 125:1–6.
- Rauch SL, Shin LM, Segal E, Pitman RK, Carson MA, McMullin K, et al (2003): Selectively reduced regional cortical volumes in post-traumatic stress disorder. *Neuroreport* 14:913–916.
- Rauch SL, van der Kolk BA, Fiesler RE, Alpert NM, Orr SP, Savage CR, et al (1996): A symptom provocation study of posttraumatic stress disorder using positron emission tomography and script-driven imagery. *Arch Gen Psychiatry* 53:380–387.
- Raz N, Gunning-Dixon F, Head D, Rodrigue KM, Williamson A, Acker JD (2004): Aging, sexual dimorphism, and hemispheric asymmetry of the cerebral cortex: Replicability of regional differences in volume. *Neurobiol Aging* 25:377–396.
- Raz N, Gunning-Dixon FM, Head D, Dupuis JH, Acker JD (1998): Neuroanatomical correlates of cognitive aging: Evidence from structural magnetic resonance imaging. *Neuropsychology* 12:95–114.
- Raz N, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, et al (1997): Selective aging of the human cerebral cortex observed in vivo: Differential vulnerability of the prefrontal gray matter. *Cereb Cortex* 7:268–282.
- Reiss AL, Hennessey JG, Rubin M, Beach L, Abrams MT, Warsofsky IS, et al (1998): Reliability and validity of an algorithm for fuzzy tissue segmentation of MRI. *J Comput Assist Tomogr* 22:471–479.
- Sanchez MM, Young LJ, Plotsky PM, Insel TR (1999): Autoradiographic and in situ hybridization localization of corticotropin-releasing factor 1 and 2 receptors in nonhuman primate brain. *J Comp Neurol* 408: 365–377.
- Selzer ML (1971): The Michigan Alcoholism Screening Test: The quest for a new diagnostic instrument. *Am J Psychiatry* 127:1653–1658.
- Shin LM, Kosslyn SM, McNally RJ, Alpert NM, Thompson WL, Rauch SL, et al (1997): Visual imagery and perception in posttraumatic stress disorder. A positron emission tomographic investigation. *Arch Gen Psychiatry* 54: 233–241.
- Shin LM, McNally RJ, Kosslyn SM, Thompson WL, Rauch SL, Alpert NM, et al (1999): Regional cerebral blood flow during script-driven imagery in childhood sexual abuse-related PTSD: A PET investigation. *Am J Psychiatry* 156:575–584.
- Shin LM, Whalen PJ, Pitman RK, Bush G, Macklin ML, Lasko NB, et al (2001): An fMRI study of anterior cingulate function in posttraumatic stress disorder. *Biol Psychiatry* 50:932–942.
- Taber KH, Rauch SL, Lanius RA, Hurley RA (2003): Functional magnetic resonance imaging: Application to posttraumatic stress disorder. *J Neuropsychiatry Clin Neurosci* 15:125–129.
- Talairach J, Tournoux P (1988): *Co-Planar Stereotaxic Atlas of the Human Brain*. New York: Thieme Medical.
- Villareal G, King CY (2001): Brain imaging in posttraumatic stress disorder. *Semin Clin Neuropsychiatry* 6:131–145.
- Vogt BA, Finch DM, Olson CR (1992): Functional heterogeneity in cingulate cortex: The anterior executive and posterior evaluative regions. *Cereb Cortex* 2:435–443.
- Vogt BA, Nimchinsky EA, Vogt LJ, Hof PR (1995): Human cingulate cortex: Surface features, flat maps, and cytoarchitecture. *J Comp Neurol* 359: 490–506.
- Vogt BA, Pandya DN (1987): Cingulate cortex of the rhesus monkey: II. Cortical afferents. *J Comp Neurol* 262:271–289.
- Vogt BA, Pandya DN, Rosene DL (1987): Cingulate cortex of the rhesus monkey: I. Cytoarchitecture and thalamic afferents. *J Comp Neurol* 262: 256–270.
- Vogt BA, Rosene DL, Pandya DN (1979): Thalamic and cortical afferents differentiate anterior from posterior cingulate cortex in the monkey. *Science* 204:205–207.
- Wellman CL (2001): Dendritic reorganization in pyramidal neurons in medial prefrontal cortex after chronic corticosterone administration. *J Neurobiol* 49:245–253.
- Yamasue H, Kasai K, Iwanami A, Ohtani T, Yamada H, Abe O, et al (2003): Voxel-based analysis of MRI reveals anterior cingulate gray-matter volume reduction in posttraumatic stress disorder due to terrorism. *Proc Natl Acad Sci U S A* 100:9039–9043.
- Zubieta JK, Chinitz JA, Lombardi U, Fig LM, Cameron OG, Liberzon I (1999): Medial frontal cortex involvement in PTSD symptoms: A SPECT study. *J Psychiatry Res* 33:259–264.